

CROSSTALK

CrossTalk proposal: Rotors have been demonstrated to drive human atrial fibrillationSanjiv M. Narayan^{1,2} and José Jalife³¹Department of Medicine/Cardiology, University of California, San Diego, CA, USA²Veterans Affairs Medical Center, San Diego, CA, USA³Center for Arrhythmia Research and Department of Internal Medicine/Cardiovascular Medicine, University of Michigan, Ann Arbor, MI, USA

Email: jjalife@umich.edu

Introduction

It is our pleasure to debate this important and clinically relevant topic. Atrial fibrillation (AF) is a leading cause of hospitalization and death (Calkins *et al.* 2012), and the need to improve AF therapy is urgent and requires dramatic advances in our mechanistic understanding. Decades of research have not dented the incidence of AF, now at epidemic proportions (Calkins *et al.* 2012), nor reduced its morbidity and mortality with the laudable exception of antithrombotic therapy (Calkins *et al.* 2012).

Our position

AF is initiated by triggers from regions including the pulmonary veins (Calkins *et al.* 2012), then sustained by specific mechanisms. We will present substantial evidence that these sustaining mechanisms include localized rotors in many patients.

Our position that 'rotors have been demonstrated to drive [human] AF'

summarizes a century of bench-to-bedside studies, culminating in recent multicentre clinical trials in which ablation of AF rotors has eliminated AF and improved patient outcomes. Conversely, there is a paucity of evidence supporting the opposing notion: that disorganized activity *per se* sustains AF without underlying driving mechanisms.

Many studies that propose disorganized mechanisms are limited to 'long-standing persistent AF'. Conversely, we provide evidence for a comprehensive mechanistic framework in which rotors sustain AF across clinical 'classes'. The terms paroxysmal AF (self-limiting), persistent and longstanding persistent AF (continuous for >7 days and >1 year, respectively) are highly dependent on how often ECGs are obtained, since AF is often asymptomatic (Charitos *et al.* 2012) and patients with persistent AF may spontaneously present in sinus rhythm (Calkins *et al.* 2012) and could thus be reclassified as paroxysmal AF (Jahangir *et al.* 2007).

Definitions

While AF appears disorganized on the ECG, the atria clearly exhibit regions of spatio-temporal organization and disorganization. The debate on whether disorganization in AF is *caused* ('driven') by sources (Pandit & Jalife, 2013) or whether disorganization *per se* sustains AF (Fig. 1) has been ongoing for a century (Jalife, 2011).

Spatially localized sources can be divided into rotors and focal sources. We define a 'rotor' as the phase singularity whose reverberations radiate 'spiral waves' at high speed into surrounding tissue (Pandit & Jalife, 2013). The spatial domain of 1:1 activation by a rotor depends on its

frequency relative to the heterogeneity of local refractory periods. In the context of cardiac arrhythmias, rotors are highly localized drivers or organizing sources of reentrant tachycardia and fibrillation (Pandit & Jalife, 2013). Although not our primary topic, a focal source is an ectopic site from where activation spreads centrifugally based on regional refractoriness and conduction (Pandit & Jalife, 2013). Rotors have long been demonstrated in animal models of AF (Pandit & Jalife, 2013), and more recently in human AF where brief targeted ablation alone can terminate AF (Narayan *et al.* 2013b) and eliminate AF long term (Narayan *et al.* 2013a).

Conversely, it has been posited that AF depends on the disorganization of wavelets, itself due to longitudinal/transmural dissociation (de Groot *et al.* 2010). However, this is supported by mapping <10–20% of MRI-quantified atrial areas (Jadidi *et al.* 2013) in small numbers of patients without proof of causality (de Groot *et al.* 2010).

Evidence for our position

Many diverse observations are readily explained and in fact predicted by our position, yet unexplained by disorganization alone (Fig. 1). Human AF resistant even to electrical cardioversion can be terminated by very few or even a single (Herweg *et al.* 2003) ablation lesion(s) (Calkins *et al.* 2012; Narayan *et al.* 2012b), or by localized physical pressure (Tzou *et al.* 2011). Propagation vectors in AF repeat reproducibly over time (Gerstenfeld *et al.* 1992) and exhibit consistent spatio-temporal features (Sahadevan *et al.* 2004; Sanders *et al.* 2005; Aienza *et al.* 2009). These observations

Sanjiv M. Narayan is Professor of Medicine at the University of California San Diego, where he treats patients with heart rhythm disorders. He was trained in software engineering and neuroscience in addition to cardiology and clinical electrophysiology. He directs an actively funded translational laboratory that uses bioengineering solutions to understand arrhythmia mechanisms, and has pioneered unique therapies for cardiac fibrillation. He has authored or co-authored over 152 original papers and review articles. **José Jalife** is the Cyrus and Jane Farrehi Professor of Cardiovascular Research and Professor of Internal Medicine at the University of Michigan. He is a leader in the study of mechanisms of cardiac arrhythmias. His work has led to major advances toward elucidating the molecular and cellular bases of atrial fibrillation, ventricular fibrillation and sudden cardiac death. He has published more than 300 original papers and review articles, and has edited/authored 15 books, including the internationally acclaimed *Cardiac Electrophysiology: From Cell to Bedside*, now in its sixth edition.



are difficult to reconcile with disorganized sustaining mechanisms, yet support spatially localized AF-sustaining regions in these patients.

Direct evidence for localized AF-sustaining mechanisms, including rotors, has been established in a variety of experimental systems. Optical mapping that utilizes voltage-sensitive dyes and contemporary signal-processing algorithms (Gray *et al.* 1998) has demonstrated AF rotors producing spatial gradients in the distribution of dominant atrial frequencies, the highest frequency lying at the rotor location (Fig. 1; Mansour *et al.* 2001). Clinical interventions show that ablation of high dominant frequency sites that eliminates left-to-right atrial frequency gradients predicts long-term freedom from AF (Atienza *et al.* 2009).

Direct clinical evidence for rotors now exists in patients with paroxysmal, persistent and longstanding persistent AF. Since human AF is spatially non-uniform, we used bi-atrial basket catheters (Narayan *et al.* 2013b) to simultaneously map much larger areas than in previous studies (de Groot *et al.* 2010) and then applied phase analysis to detect regions that activate sequentially but may be obscured by the fibrillatory milieu. Simultaneous analysis of wide areas increases the sensitivity to detect rotors that precess (show limited meander), and is essential to prove or disprove their existence. Yet, to the best of our knowledge, simultaneous analysis of wide areas is absent in studies supporting disorganized mechanisms (de Groot *et al.* 2010; Lee *et al.* 2014). Although higher

spatial resolution would be welcome, we reasoned from minimum bi-atrial (Narayan *et al.* 2012a) repolarization (~ 110 ms) and conduction velocity (~ 40 cm s⁻¹) that human atria can support 1:1 conduction from a rotor in circuit paths as short as ~ 4 –5 cm (Rensma *et al.* 1988). Clinical electrode spacing can map such reentry (Rappel & Narayan, 2013).

The CONFIRM trial (CONventional ablation for AF with or without Focal Impulse and Rotor Modulation, FIRM) used this approach in 92 patients at 107 procedures to reveal rotors (or focal sources) in 97% of patients with paroxysmal, persistent and longstanding persistent AF. Rotors were concurrent (2.1 ± 1.0 per patient), more prevalent with advanced AF (Narayan *et al.* 2012b), and precessed in ~ 2 –3 cm² areas (Narayan *et al.* 2013b) causing irregular electrograms at the rotor tip (Zlochiver *et al.* 2008) and varying spiral waves from fibrillatory disorganization. Notably, causality was proven by the ability of brief targeted ablation to eliminate AF acutely and on long-term follow-up, with (Narayan *et al.* 2012b) or without (Narayan *et al.* 2013a) other ablation. These results are now independently validated (Shivkumar *et al.* 2012; Miller *et al.* 2013), and laboratories worldwide have identified rotors in human AF using diverse methods, including phase mapping (Ghoraani *et al.* 2013; Haissaguerre *et al.* 2013; Lin *et al.* 2013; Lee *et al.* 2014). Studies are defining how localized ablation eliminates AF, possibly analogous to the proven elimination of micro-reentrant atrial tachycardia by focal ablation.

Limitations of the opposing position

While it is often assumed that Maze surgery supports disorganized mechanisms (Moe *et al.* 1964), the numbers of lesions (Cox *et al.* 1991) are insufficient to constrain dozens or hundreds of the wavelets proposed (de Groot *et al.* 2010). Since rotors need 'elbow room' (typically $2 \times$ the centre of rotation or ~ 5 –10 mm), Maze may in fact simply prevent rotors from sustaining. Indeed, one original motivation for Maze was to intersect relatively large reentrant drivers (Cox *et al.* 1991).

Notably, studies supporting AF disorganization involved few patients ($n = 49$) (de Groot *et al.* 2010), and mapped only 10–21 cm² or <10 –20% of MRI-defined left atrial areas (100–138 cm²) (Jadidi *et al.* 2013). It is unclear why this work (de Groot *et al.* 2010) consistently fails to show any rotational activity in human AF, since others now show rotations of varying duration depending on the size of the mapping plaque and technique (Ghoraani *et al.* 2013; Haissaguerre *et al.* 2013; Lin *et al.* 2013; Narayan *et al.* 2013b; Lee *et al.* 2014). It is possible that activation mapping of these difficult-to-interpret AF electrograms (Fig. 1; de Groot *et al.* 2010) obscures rotation evident in other approaches (e.g. phase mapping). Importantly, to the best of our knowledge, interventions have never been used to prove that disorganization is an AF-sustaining mechanism rather than just a bystander to another mechanism (e.g. rotors). Finally, the disorganized AF hypothesis is inconsistent with and cannot easily explain the wealth of clinical and experimental data supporting localized AF sources in at least some patients as discussed above.




Figure 1. Lines of evidence favouring rotors over disorganized mechanisms for AF

Left human rotor (where ablation terminated AF) surrounded by fibrillatory conduction. Right, disorganized AF maps, but created from analysis of difficult-to-interpret AF electrograms (de Groot *et al.* 2010).

Conclusions and future directions

Rotors have been proven to sustain AF in patients in all clinical subtypes and numerous animal models. These data come from various groups using diverse methods with interventions and long-term outcomes to prove causality. The opposing contention, that rotors do not exist but that AF is sustained by disorganization without driving sources, is supported only circumstantially by experiments with important technical limitations, in a small numbers of patients, and with no proof of causality. Mapping and therapy of AF rotors has already substantially improved clinical outcomes in multicentre trials. Future work should define how electrical, structural or neural remodelling contributes to the formation of AF sources and disorganization from them to remaining tissue.

Call for comments

Readers are invited to give their views on this and the accompanying CrossTalk articles in this issue by submitting a brief comment. Comments may be posted up to 6 weeks after publication of the article, at which point the discussion will close and authors will be invited to submit a 'final word'. To submit a comment, go to <http://jp.physoc.org/letters/submit/jphysiol;592/15/3163>

References

- Atienza F, Almendral J, Jalife J, Zlochiver S, Ploutz-Snyder R, Torrecilla EG, Arenal A, Kalifa J, Fernandez-Aviles F & Berenfeld O (2009). Real-time dominant frequency mapping and ablation of dominant frequency sites in atrial fibrillation with left-to-right frequency gradients predicts long-term maintenance of sinus rhythm. *Heart Rhythm* **6**, 33–40.
- Calkins H, Kuck KH, Cappato R, Brugada J, Camm AJ, Chen SA, Crijns HJ, Damiano RJ Jr, Davies DW, DiMarco J, *et al.* (2012). 2012 HRS/EHRA/ECAS expert consensus statement on catheter and surgical ablation of atrial fibrillation: recommendations for patient selection, procedural techniques, patient management and follow-up, definitions, endpoints, and research trial design. *Heart Rhythm* **9**, 632–696.
- Charitos EI, Stierle U, Ziegler PD, Baldewig M, Robinson DR, Sievers HH & Hanke T (2012). A comprehensive evaluation of rhythm monitoring strategies for the detection of atrial fibrillation recurrence: insights from 647 continuously monitored patients and implications for monitoring after therapeutic interventions. *Circulation* **126**, 806–814.
- Cox JL, Canavan TE, Schuessler RB, Cain ME, Lindsay BD, Stone C, Smith PK, Corr PB & Boineau JP (1991). The surgical treatment of atrial fibrillation. II. Intraoperative electrophysiologic mapping and description of the electrophysiologic basis of atrial flutter and atrial fibrillation. *J Thorac Cardiovasc Surg* **101**, 406–426.
- de Groot NM, Houben RP, Smeets JL, Boersma E, Schotten U, Schalij MJ, Crijns H & Allessie MA (2010). Electrophathological substrate of longstanding persistent atrial fibrillation in patients with structural heart disease: epicardial breakthrough. *Circulation* **122**, 1674–1682.
- Gerstenfeld E, Sahakian A & Swiryn S (1992). Evidence for transient linking of atrial excitation during atrial fibrillation in humans. *Circulation* **86**, 375–382.
- Gharaani B, Dalvi R, Gizurason S, Das M, Ha A, Suszko A, Krishnan S & Chauhan VS (2013). Localized rotational activation in the left atrium during human atrial fibrillation: Relationship to complex fractionated atrial electrograms and low-voltage zones. *Heart Rhythm* **10**, 1830–1838.
- Gray RA, Pertsov AM & Jalife J (1998). Spatial and temporal organization during cardiac fibrillation. *Nature* **392**, 75–78.
- Haissaguerre M, Hocini M, Shah AJ, Derval N, Sacher F, Jais P & Dubois R (2013). Noninvasive panoramic mapping of human atrial fibrillation mechanisms: a feasibility report. *J Cardiovasc Electrophysiol* **24**, 711–717.
- Herweg B, Kowalski M & Steinberg JS (2003). Termination of persistent atrial fibrillation resistant to cardioversion by a single radiofrequency application. *Pacing Clin Electrophysiol* **26**, 1420–1423.
- Jadidi AS, Cochet H, Shah AJ, Kim SJ, Duncan E, Miyazaki S, Serresant M, Lehmann H, Lederlin M, Linton N, Forclaz A, Nault I, Rivard L, Wright M, Liu X, Scherr D, Wilton SB, Roten L, Pascale P, Derval N, Sacher F, Knecht S, Keyl C, Hocini M, Montaudon M, Laurent F, Haissaguerre M & Jais P (2013). Inverse relationship between fractionated electrograms and atrial fibrosis in persistent atrial fibrillation: combined magnetic resonance imaging and high-density mapping. *J Am Coll Cardiol* **62**, 802–812.
- Jahangir A, Lee V, Friedman PA, Trusty J, Hodge D, Kopecky S, Packer DL, Hammill SC, Shen W & Gersh BJ (2007). Long-term progression and outcomes with aging in patients with lone atrial fibrillation: a 30-year follow-up study. *Circulation* **115**, 3050–3056.
- Jalife J (2011). Déjà vu in the theories of atrial fibrillation dynamics. *Cardiovasc Res* **89**, 766–775.
- Lee G, Kumar S, Teh A, Madry A, Spence S, Larobina M, Goldblatt J, Brown R, Atkinson V, Moten S, Morton JB, Kistler PM, Sanders P & Kalman JM (2014). Epicardial wave mapping in human long-lasting persistent atrial fibrillation: transient rotational circuits, complex wavefronts, and disorganized activity. *Eur Heart J* **35**, 86–97.
- Lin YJ, Lo MT, Lin C, Chang SL, Lo LW, Hu YF, Hsieh WH, Chang HY, Lin WY, Chung FP, Liao JN, Chen YY, Hanafy D, Huang NE & Chen SA (2013). Prevalence, characteristics, mapping, and catheter ablation of potential rotors in nonparoxysmal atrial fibrillation. *Circ Arrhythm Electrophysiol* **6**, 851–858.
- Mansour M, Mandapati R, Berenfeld O, Chen J, Samie FH & Jalife J (2001). Left-to-right gradient of atrial frequencies during acute atrial fibrillation in the isolated sheep heart. *Circulation* **103**, 2631–2636.
- Miller JM, Krummen DE, Clopton P, Day JD, Daubert JP, Ellenbogen KA, Hummel JD, Kowal RC, Mansour MC, Reddy VY, Shivkumar K, Steinberg JS, Swarup V, Wheelan KR & Narayan SM (2013). Ablation of Atrial Fibrillation Rotors and Focal Sources Improves Outcome Over Conventional Ablation Alone in Independent Laboratories: Multicenter Validation of Focal Impulse And Rotor Modulation (FIRM). *Circulation* **128**, A16890.
- Moe GK, Rheinboldt W & Abildskov J (1964). A computer model of atrial fibrillation. *Am Heart J* **67**, 200–220.
- Narayan SM, Krummen DE, Donsky A, Swarup V & Miller JM (2013a). Precise rotor elimination without concomitant pulmonary vein isolation for the successful elimination of paroxysmal atrial fibrillation. *PRECISE-PAF. Heart Rhythm* **10**, LBCT4.
- Narayan SM, Krummen DE & Rappel W-J (2012a). Clinical mapping approach to identify rotors and focal beats in human atrial fibrillation. *J Cardiovasc Electrophysiol* **23**, 447–454.
- Narayan SM, Krummen DE, Shivkumar K, Clopton P, Rappel W-J & Miller J (2012b). Treatment of atrial fibrillation by the ablation of localized sources: CONFIRM (conventional ablation for atrial fibrillation with or without focal impulse and rotor modulation) trial. *J Am Coll Cardiol* **60**, 628–636.
- Narayan SM, Shivkumar K, Krummen DE, Miller JM & Rappel W-J (2013b). Panoramic electrophysiological mapping but not electrogram morphology identifies stable sources for human atrial fibrillation. Stable atrial fibrillation rotors and focal sources relate poorly to fractionated electrograms. *Circ Arrhythm Electrophysiol* **6**, 58–67.
- Pandit SV & Jalife J (2013). Rotors and the dynamics of cardiac fibrillation. *Circ Res* **112**, 849–862.

- Rappel W-J & Narayan SM (2013). Theoretical considerations for mapping activation in human cardiac fibrillation. *Chaos* **23**, 023113.
- Rensma P, Allesie M, Lammers W, Bonke F & Schalij M (1988). Length of excitation wave and susceptibility to reentrant atrial arrhythmias in normal conscious dogs. *Circ Res* **62**, 395–410.
- Sahadevan J, Ryu K, Peltz L, Khrestian CM, Stewart RW, Markowitz AH & Waldo AL (2004). Epicardial mapping of chronic atrial fibrillation in patients: preliminary observations. *Circulation* **110**, 3293–3299.
- Sanders P, Berenfeld O, Hocini M, Jaïs P, Vaidyanathan R, Hsu LF, Garrigue S, Takahashi Y, Rotter M, Sacher F, Scavée C, Ploutz-Snyder R, Jalife J & Haïssaguerre M (2005). Spectral analysis identifies sites of high-frequency activity maintaining atrial fibrillation in humans. *Circulation* **112**, 789–797.
- Shivkumar K, Ellenbogen KA, Hummel JD, Miller JM & Steinberg JS (2012). Acute termination of human atrial fibrillation by identification and catheter ablation of localized rotors and sources: first multicenter experience of focal impulse and rotor modulation (FIRM) ablation. *J Cardiovasc Electrophysiol* **23**, 1277–1285.
- Tzou WS, Saghy L & Lin D (2011). Termination of persistent atrial fibrillation during left atrial mapping. *J Cardiovasc Electrophysiol* **22**, 1171–1173.
- Zlochiver S, Yamazaki M, Kalifa J & Berenfeld O (2008). Rotor meandering contributes to irregularity in electrograms during atrial fibrillation. *Heart Rhythm* **5**, 846–854.

Additional information

Competing interests

S. M. Narayan is co-author of intellectual property owned by the University of California Regents and licensed to Topera Inc. (San Diego, CA, USA). Topera does not sponsor any research, including that presented here. S. M. Narayan holds equity in Topera, and reports having received honoraria from Medtronic, St Jude Medical and Biotronik. J. Jalife serves on the Scientific Advisory Board of Topera, Inc.

Funding

The authors' work is supported by National Heart, Lung, and Blood Institute grants HL103800 and HL83559 (S.M.N.) and the Leducq Foundation (J.J.).